

- Bormans" (Ann. & Mag. Nat. Hist. (7) xi. pp. 231-241 & 266-270, 1903).
- BURN, MALCOLM. (1904²). "Observations on the Dermaptera, including Revisions of several Genera and Descriptions of New Species" (Tr. Ent. Soc. London, pp. 277-322, 1904).
- (1907¹). "A Preliminary Revision of the Forficulidæ (sensu stricto) and of the Chelisochidæ, Families of the Dermaptera" (Tr. Ent. Soc. Lond. pp. 91-134, pl. iv., 1907).
- (1907²). "Ueber einige neue und interessante Dermapteren Arten aus Kamerun und Togo" (Deutsch. ent. Zeitschr. p. 487, 1907).
- (1908¹). "Notes on the Forficularia.—XI. On new and little-known Species and Synonymic Notes" (Ann. & Mag. Nat. Hist. (8) i. pp. 47-51, 1908).
- (1908²). "Two new Dermaptera in the Collection of the Leyden Museum" (Notes from the Leyden Museum, xxx., Note xiii. pp. 95 & 96, 1908).
- (1908³). "Sopra alcuni Dermatteri del Museo Civico di Genova" (Bull. Soc. Ent. Ital. ix. 1908, pp. 175-181).
- GERSTAECKER. (1873). "Beitrag zur Kenntniss der Orthopteren-Fauna Guineas nach den von T. Buchholz während den Jahren 1872 bis 1875 daselbst gesammelten Arten" (Mitth. des naturwissenschaftl. Vereins für Neu-vorpommern und Rügen, 1883).
- HAAN, DR. W. DE. (1842). "Bijdragen tot de Kennis der Orthoptera" (Verhandl. over de natuurlijke Geschiedenis der Nederlandsche overzeesche Bezittingen, 1842).
- KARSCH, DR. F. (1886). "Ueber einige neue oder weniger bekannte Ohrwuermer (Dermaptera) der äthiopischen Region" (Berliner ent. Zeitschr. Bd. xxx. Heft 1, p. 85, 1886).
- KIRBY, W. F. (1891). "A Revision of the Forficulidæ, with Descriptions of New Species in the British Museum" (Journ. Linn. Soc. vol. xxiii. pp. 502-531, pl. xii., 1891).
- (1904). Syn. Cat. Orth. i. London (1904).
- REHN, JAMES A. G. (1903²). "Studies in American Forficulidæ" (Proc. Nat. Sci. Phil. p. 209, 1903).
- SCUDDER, SAMUEL H. (1869). Proc. Bost. Soc. N. H. xii. p. 344.
- SERVILLE, AUDINET. (1839). 'Suite à Buffon. Histoire Naturelle des Insectes Orthoptères.' Paris, 1839.
- STAL, C. (1855). "Entomologiska Notiser" (O. V. A. F. xii. 1855, p. 347).

XV.—*On some new Species of Birds from Katanga, Congo Free State.* By S. A. NEAVE, M.A., B.Sc.

I HAVE found the following apparently new species among the birds in my collection made during 1907 in the Katanga region of the Congo Free State.

Trochocercus vivax, sp. n.

T. similis T. livittata, sed plaga alari alba nulla; corporis lateribus griseescentibus, pectore et abdomine albis; rostro cyanescenti-

corneo, apice fusciscente; pedibus cyaescenti-griscis; iride brunnea.

Long. tot. c. 6.1 poll., culm. 0.7, alæ 2.75, caudæ 3.2, tarsi 0.75.

Hab. Katanga, S.E. Congo Free State.

Criniger sylvicultor, sp. n.

C. similis C. cabanisi, sed supra pallidior, gutture et pectore toto pallide sulfureo-flavis; præpectore olivascento sulfureo lavato; rostro fusciscenti-corneo, subtus pallidior; pedibus pallide griscentibus; iride olivascenti-griseo.

Long. tot. c. 7.0 poll., culm. 0.85, alæ 3.65, caudæ 3.6, tarsi 1.05.

Hab. Katanga.

Calamonastes katangæ, sp. n.

C. similis C. cinereo, sed brunnescentior; pectore et abdomino albidis; gula albida, fusco variegato; gutture summo et præpectore cineraceo, fasciam latam formantibus; rostro nigro; pedibus flavescenti-carneis; iride rufescenti-brunneo.

Long. tot. c. 4.1 poll., culm. 0.55, alæ 2.3, caudæ 1.7, tarsi 0.8.

Hab. Katanga.

Dryodromas pearsoni, sp. n.

Similis *D. rufifronti* et cauda nigra; reatricibus ad apicem cinereis, pileo ferrugineo; dorso rufescenti-brunneo; uropygio et supracaudalibus cinerascensibus; subtus alba, gutture et præpectore pallide cervino tinctis; hypochondriis cinereis distinguenda; rostro brunnescenti-corneo, mandibula pallide carnea; pedibus brunnescenti-carneis; iride flavicanti-brunnea.

Long. tot. c. 4.1 poll., culm. 0.45, alæ 2.0, caudæ 2.0, tarsi 0.65.

Hab. Lufupa River, Western Katanga.

Dedicated to Dr. Arthur Pearson, Principal Medical Officer to the Tanganyika Concession Ltd., Kambore.

XVI.—*Descriptions of new Genera and Species of New-Zealand Coleoptera.* By Major T. BROUN, F.E.S.

[Continued from p. 71.]

Group *Erirhinidæ*.

Erirhinus insignis, sp. n.

Subovate, moderately elongate, nitid, sparsely clothed with fine yellowish hairs, which are somewhat concentrated on